

Combinatorics of symbolic Rees algebras of edge ideals of clutters

José Martínez-Bernal, Carlos Rentería-Márquez, and Rafael H. Villarreal

This paper is dedicated to Wolmer Vasconcelos.

ABSTRACT. Let $\mathcal{C}$ be a clutter and let I be its edge ideal. We present a combinatorial description of the minimal generators of the symbolic Rees algebra $R_s(I)$ of I . It is shown that the minimal generators of $R_s(I)$ are in one to one correspondence with the indecomposable parallelizations of $\mathcal{C}$. From our description some major results on symbolic Rees algebras of perfect graphs and clutters will follow. As a byproduct, we give a method, using Hilbert bases, to compute all indecomposable parallelizations of $\mathcal{C}$ and all the corresponding vertex covering numbers.

1. Introduction

Let $\mathcal{C}$ be a *clutter* with finite vertex set $X = \{x_1, \dots, x_n\}$, i.e., $\mathcal{C}$ is a family of subsets of X , called edges, none of which is included in another. The set of vertices and edges of $\mathcal{C}$ are denoted by $V(\mathcal{C})$ and $E(\mathcal{C})$ respectively. A basic example of a clutter is a graph. Let $R = K[x_1, \dots, x_n]$ be a polynomial ring over a field K . The *edge ideal* of $\mathcal{C}$, denoted by $I = I(\mathcal{C})$, is the ideal of R generated by all square-free monomials $x_e = \prod_{x_i \in e} x_i$ such that $e \in E(\mathcal{C})$. The assignment $\mathcal{C} \mapsto I(\mathcal{C})$ gives a natural one to one correspondence between the family of clutters and the family of square-free monomial ideals.

The *blowup algebra* studied here is the *symbolic Rees algebra* of I :

$$R_s(I) = R \oplus I^{(1)}t \oplus \dots \oplus I^{(i)}t^i \oplus \dots \subset R[t],$$

where t is a new variable and $I^{(i)}$ is the i th symbolic power of I . Recall that the i th symbolic power of I is defined as

$$I^{(i)} = S^{-1}I^i \cap R,$$

where $S = R \setminus \cup_{k=1}^s \mathfrak{p}_k$, the ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_s$ are the minimal primes of I and $S^{-1}I^i$ is the localization of I^i at S . In our situation the i th symbolic power of I can be

2000 *Mathematics Subject Classification.* Primary 13A30; Secondary 13F20, 05C65, 05C75.

The second author was Partially supported by COFAA-IPN and SNI. The third author was partially supported by CONACyT grant 49251-F and SNI.

expressed using systems of linear inequalities (see Eq. (2.1) in Section 2). Closely related to $R_s(I)$ is—another blowup algebra—the *Rees algebra* of I :

$$R[It] = R \oplus It \oplus \cdots \oplus I^i t^i \oplus \cdots = K[\{x_1, \dots, x_n, x_e t \mid e \in E(\mathcal{C})\}] \subset R[t].$$

Blowup algebras are interesting objects of study in algebra and geometry [30].

The study of symbolic powers of edge ideals from the point of view of graph theory and combinatorics was initiated in [28] and further elaborated on in [29, 32]. A breakthrough in this area is the translation of combinatorial problems (e.g., the Conforti-Cornuéjols conjecture [6], the max-flow min-cut property, the idealness of a clutter, or the integer rounding property) into algebraic problems of blowup algebras of edge ideals [3, 8, 10, 11, 16, 17, 22].

By a result of Lyubeznik [21], $R_s(I)$ is a K -algebra of finite type generated by a unique minimal finite set of monomials. The main theorem of this paper is a description—in combinatorial optimization terms—of this minimal set of generators of $R_s(I)$ as a K -algebra. Before stating the theorem, we need to recall some more terminology and notations.

A subset C of X is called a *vertex cover* of $\mathcal{C}$ if every edge of $\mathcal{C}$ contains at least one vertex of C . A vertex cover C is called a *minimal vertex cover* if no proper subset of C is a vertex cover. The number of vertices in a minimum vertex cover of $\mathcal{C}$, denoted by $\alpha_0(\mathcal{C})$, is called the *vertex covering number* of $\mathcal{C}$. The dual concept of a vertex cover is a *stable set*, i.e., a subset C of X is a vertex cover of $\mathcal{C}$ if and only if $X \setminus C$ is a stable set. The number of vertices in a maximum stable set, denoted by $\beta_0(\mathcal{C})$, is called the *stability number* of $\mathcal{C}$. Notice that $\alpha_0(\mathcal{C}) + \beta_0(\mathcal{C}) = n$.

A clutter $\mathcal{C}$ is called *indecomposable* if it cannot be decomposed as a disjoint union of induced subclutters $\mathcal{C}_1, \mathcal{C}_2$ such that $\alpha_0(\mathcal{C}) = \alpha_0(\mathcal{C}_1) + \alpha_0(\mathcal{C}_2)$ (see Definition 2.2). Erdős and Gallai [13] introduced this notion for graphs. A clutter obtained from $\mathcal{C}$ by a sequence of deletions and duplications of vertices is called a *parallelization* (see Definition 2.3). If $a = (a_i)$ is a vector in $\mathbb{N}^n$, we denote by $\mathcal{C}^a$ the clutter obtained from $\mathcal{C}$ by successively deleting any vertex x_i with $a_i = 0$ and duplicating $a_i - 1$ times any vertex x_i if $a_i \geq 1$ (see Example 2.4).

Our main result is:

Theorem 2.6 *Let $0 \neq a = (a_i) \in \mathbb{N}^n$, $b \in \mathbb{N}$. Then $x_1^{a_1} \cdots x_n^{a_n} t^b$ is a minimal generator of $R_s(I)$, as a K -algebra, if and only if $\mathcal{C}^a$ is an indecomposable clutter and $b = \alpha_0(\mathcal{C}^a)$.*

There are two cases where a combinatorial description of the symbolic Rees algebra is known. If the clutter $\mathcal{C}$ has the max-flow min-cut property, then by a result of [17], we have $I^i = I^{(i)}$ for all $i \geq 1$, i.e., $R_s(I) = R[It]$ and a minimal generator of $R_s(I)$ is either a vertex x_i or an “edge” $x_e t$ with $e \in E(\mathcal{C})$. If $\mathcal{C}$ is a perfect graph, then the minimal generators of $R_s(I)$ are in one to one correspondence with the cliques (complete subgraphs) of $\mathcal{C}$ [32]. Both cases will follow from our combinatorial description of $R_s(I)$ (see Corollaries 2.10 and 4.2 respectively).

As a byproduct, in Section 3 we give a method—based on the computation of Hilbert bases of polyhedral cones—to compute all indecomposable parallelizations of any clutter $\mathcal{C}$ along with all the corresponding vertex covering numbers. In particular our method allows to compute all indecomposable induced subclutters of any clutter $\mathcal{C}$. This means that the symbolic Rees algebra of I encodes combinatorial information of the clutter which can be decoded using a computer program, such as NORMALIZ [4], which is able to compute Hilbert bases of polyhedral cones.

Harary and Plummer [19] studied some properties of indecomposable graphs. They showed that if a connected graph is separated by the points of a complete subgraph, then G is decomposable. All indecomposable graphs with at least three vertices contain at least one odd cycle, and the join of two indecomposable graphs is indecomposable [19]. Indecomposable graphs were first studied from an algebraic point of view in [12]. To the best of our knowledge there is no structure theorem for indecomposable graphs.

Indecomposable subgraphs occur naturally in the theory of perfect graphs. Indeed, a graph G is perfect if and only if the indecomposable parallelizations of G are exactly the complete subgraphs or cliques of G (see Proposition 4.1). This was first shown in [12] using the main result of [5]. For graphs, we can use our methods to locate all *induced odd cycles* (*odd holes*) and all *induced complements of odd cycles* (*odd antiholes*) of length at least five. Indeed, odd holes of any length and odd antiholes of length at least five are indecomposable subgraphs (see Lemma 3.5), and thus by Theorem 2.6 they correspond to minimal generators of the symbolic Rees algebra of the edge ideal of the graph. Odd holes and odd antiholes play a major role in graph theory. In [5] it is shown that a graph G is perfect if and only if G is a *Berge graph*, i.e., if and only if G has no odd holes or odd antiholes of length at least five. In commutative algebra odd holes occurred for the first time in [26], and later in the description of $I(G)^{\{2\}}$, the join of an edge ideal of a graph G with itself [27]. They also occurred in the description of the associated primes of powers of ideals of vertex covers of graphs [15].

The problem of finding a minimum vertex cover of a graph is a classical optimization problem in computer science and is a typical example of an NP-hard problem. From the point of view of computational complexity theory, finding all indecomposable subgraphs of a given graph using our method is a hard problem because to apply this method we must know all minimal vertex covers (see Section 3). Thus, although our results provide some tools for computing, the contributions of this paper could be more interesting from the theoretical point of view.

Throughout the paper we introduce most of the notions that are relevant for our purposes. For unexplained terminology, we refer to [7, 25, 30, 31].

2. Symbolic Rees algebras of edge ideals

In this section we will give a combinatorial description of the minimal generators of the symbolic Rees algebra of the edge ideal of a clutter using the notion of a parallelization of a clutter and the notion of an indecomposable clutter. We continue using the definitions and terms from the introduction.

Let $\mathcal{C}$ be a clutter with vertex set $X = \{x_1, \dots, x_n\}$ and let $I = I(\mathcal{C})$ be its edge ideal. We denote by $\Upsilon(\mathcal{C})$ the clutter whose edges are the minimal vertex covers of $\mathcal{C}$. The clutter $\Upsilon(\mathcal{C})$ is called the *blocker* of $\mathcal{C}$ or the *Alexander dual* of $\mathcal{C}$. As usual, we use x^a as an abbreviation for $x_1^{a_1} \cdots x_n^{a_n}$, where $a = (a_i) \in \mathbb{N}^n$.

If C is a subset of X , its *characteristic vector* is the vector $v = \sum_{x_i \in C} e_i$, where e_i is the i th unit vector in $\mathbb{R}^n$. Let $C_1, \dots, C_s$ be the minimal vertex covers of $\mathcal{C}$ and let u_k be the characteristic vector of C_k for $1 \leq k \leq s$. In our situation, according to [31, Proposition 7.3.14], the b th symbolic power of I has a simple expression:

$$\begin{aligned} I^{(b)} &= \mathfrak{p}_1^b \cap \cdots \cap \mathfrak{p}_s^b \\ (2.1) \quad &= (\{x^a \mid \langle a, u_k \rangle \geq b \text{ for } k = 1, \dots, s\}), \end{aligned}$$

where $\mathfrak{p}_k$ is the prime ideal of R generated by C_k and $\langle \cdot, \cdot \rangle$ denotes the standard inner product. In particular, if $b = 1$, we obtain the primary decomposition of I because $I^{(1)} = I$. Thus the height of I equals $\alpha_0(\mathcal{C})$, the vertex covering number of $\mathcal{C}$. This is a hint of the rich interaction between the combinatorics of $\mathcal{C}$ and the algebra of I .

Next, in Lemma 2.1, we give a simple description of $R_s(I)$ that was first observed in the discussion of symbolic Rees algebras given in [14, p. 75], see also [20]. Let $a = (a_i) \neq 0$ be a vector in $\mathbb{N}^n$ and let $b \in \mathbb{N}$. From Eq. (2.1) we get that $x^a t^b$ is in $R_s(I)$ if and only if

$$\langle a, u_k \rangle \geq b \text{ for } k = 1, \dots, s.$$

If a, b satisfy this system of linear inequalities, we say that a is a *b-vertex cover* of $\Upsilon(\mathcal{C})$. Often we will call a *b-vertex cover* simply a *b-cover*. Thus the symbolic Rees algebra of I is equal to the K -subalgebra of $R[t]$ generated by all monomials $x^a t^b$ such that a is a *b-cover* of $\Upsilon(\mathcal{C})$, as was first shown in [14, Theorem 3.5]. The notion of a *b-cover* occurs in combinatorial optimization (see for instance [25, Chapter 77, p. 1378] and the references there) and algebraic combinatorics [14, 20]. We say that a *b-cover* a of $\Upsilon(\mathcal{C})$ is *decomposable* if there exists an *i-cover* c and a *j-cover* d of $\Upsilon(\mathcal{C})$ such that $a = c + d$ and $b = i + j$. If a is not decomposable, we call a *indecomposable*. The indecomposable 0 and 1 covers of $\Upsilon(\mathcal{C})$ are the unit vectors $e_1, \dots, e_n$ and the characteristic vectors $v_1, \dots, v_q$ of the edges of $\mathcal{C}$ respectively.

Lemma 2.1. *A monomial $x^a t^b \neq 1$ is a minimal generator of $R_s(I)$, as a K -algebra, if and only if a is an indecomposable b -cover of $\Upsilon(\mathcal{C})$. In particular, the following equality holds:*

$$(2.2) \quad R_s(I) = K[\{x^a t^b \mid a \text{ is an indecomposable } b\text{-cover of } \Upsilon(\mathcal{C})\}].$$

PROOF. It follows from the discussion above, by decomposing any b -cover into indecomposable ones. $\square$

Let S be a set of vertices of a clutter $\mathcal{C}$. The *induced subclutter* on S , denoted by $\mathcal{C}[S]$, is the maximal subclutter of $\mathcal{C}$ with vertex set S . Thus the vertex set of $\mathcal{C}[S]$ is S and the edges of $\mathcal{C}[S]$ are exactly the edges of $\mathcal{C}$ contained in S . Notice that $\mathcal{C}[S]$ may have isolated vertices, i.e., vertices that do not belong to any edge of $\mathcal{C}[S]$. If $\mathcal{C}$ is a discrete clutter, i.e., all the vertices of $\mathcal{C}$ are isolated, we set $I(\mathcal{C}) = 0$ and $\alpha_0(\mathcal{C}) = 0$.

Let $\mathcal{C}$ be a clutter and let X_1, X_2 be a partition of $V(\mathcal{C})$ into nonempty sets. Clearly, one has the inequality

$$(2.3) \quad \alpha_0(\mathcal{C}) \geq \alpha_0(\mathcal{C}[X_1]) + \alpha_0(\mathcal{C}[X_2]).$$

If $\mathcal{C}$ is a graph and equality occurs, Erdős and Gallai [13] call $\mathcal{C}$ a *decomposable graph*. This motivates the following similar notion for clutters.

Definition 2.2. A clutter $\mathcal{C}$ is called *decomposable* if there are nonempty vertex sets X_1, X_2 such that X is the disjoint union of X_1 and X_2 , and $\alpha_0(\mathcal{C}) = \alpha_0(\mathcal{C}[X_1]) + \alpha_0(\mathcal{C}[X_2])$. If $\mathcal{C}$ is not decomposable, it is called *indecomposable*.

Examples of indecomposable graphs include complete graphs, odd cycles and complements of odd cycles of length at least five (see Lemma 3.5).

Definition 2.3. (Schrijver [25]) The *duplication* of a vertex x_i of a clutter $\mathcal{C}$ means extending its vertex set X by a new vertex x'_i and replacing $E(\mathcal{C})$ by

$$E(\mathcal{C}) \cup \{(e \setminus \{x_i\}) \cup \{x'_i\} \mid x_i \in e \in E(\mathcal{C})\}.$$

The *deletion* of x_i , denoted by $\mathcal{C} \setminus \{x_i\}$, is the clutter formed from $\mathcal{C}$ by deleting the vertex x_i and all edges containing x_i . A clutter obtained from $\mathcal{C}$ by a sequence of deletions and duplications of vertices is called a *parallelization*.

It is not difficult to verify that these two operations commute. If $a = (a_i)$ is a vector in $\mathbb{N}^n$, we denote by $\mathcal{C}^a$ the clutter obtained from $\mathcal{C}$ by successively deleting any vertex x_i with $a_i = 0$ and duplicating $a_i - 1$ times any vertex x_i if $a_i \geq 1$ (for graphs cf. [18, p. 53]).

Example 2.4. Let G be the graph whose only edge is $\{x_1, x_2\}$ and let $a = (3, 3)$. We set $x_i^1 = x_i$ for $i = 1, 2$. The parallelization G^a is a complete bipartite graph with bipartition $V_1 = \{x_1^1, x_1^2, x_1^3\}$ and $V_2 = \{x_2^1, x_2^2, x_2^3\}$. Note that x_i^k is a vertex, i.e., k is an index not an exponent.






Fig. 1. Graph

Fig. 2. Duplications of x_1 Fig. 3. Duplications of x_2

Proposition 2.5. ([9, Lemma 2.15], [25, p. 1385, Eq. (78.6)]) Let $\mathcal{C}$ be a clutter with n vertices and let $\Upsilon(\mathcal{C})$ be the blocker of $\mathcal{C}$. If $a = (a_i) \in \mathbb{N}^n$, then

$$\min \left\{ \sum_{x_i \in C} a_i \mid C \in \Upsilon(\mathcal{C}) \right\} = \alpha_0(\mathcal{C}^a).$$

We come to the main result of this section.

Theorem 2.6. Let $\mathcal{C}$ be a clutter with vertex set $X = \{x_1, \dots, x_n\}$ and let $0 \neq a = (a_i) \in \mathbb{N}^n$, $b \in \mathbb{N}$. Then $x^a t^b$ is a minimal generator of $R_s(I(\mathcal{C}))$, as a K -algebra, if and only if $\mathcal{C}^a$ is an indecomposable clutter and $b = \alpha_0(\mathcal{C}^a)$.

PROOF. We may assume that $a = (a_1, \dots, a_m, 0, \dots, 0)$, where $a_i \geq 1$ for $i = 1, \dots, m$. For each $1 \leq i \leq m$ the vertex x_i is duplicated $a_i - 1$ times, and the vertex x_i is deleted for each $i > m$. We denote the duplications of x_i by $x_i^2, \dots, x_i^{a_i}$ and set $x_i^1 = x_i$ for $1 \leq i \leq m$. Thus the vertex set of $\mathcal{C}^a$ can be written as

$$X^a = \{x_1^1, \dots, x_1^{a_1}, \dots, x_m^1, \dots, x_m^{a_m}\} = X^{a_1} \cup X^{a_2} \cup \dots \cup X^{a_m},$$

where $X^{a_i} = \{x_i^1, \dots, x_i^{a_i}\}$ for $1 \leq i \leq m$ and $X^{a_i} \cap X^{a_j} = \emptyset$ for $i \neq j$.

$\Rightarrow$) Assume that $x^a t^b$ is a minimal generator of $R_s(I(\mathcal{C}))$. Then, by Lemma 2.1, a is an indecomposable b -cover of $\Upsilon(\mathcal{C})$. First we prove that $b = \alpha_0(\mathcal{C}^a)$. There is k such that $a_k \neq 0$. We may assume that $a - e_k \neq 0$. By Proposition 2.5 we need only show the equality

$$b = \min \left\{ \sum_{x_i \in C} a_i \mid C \in \Upsilon(\mathcal{C}) \right\}.$$

As a is a b -cover of $\Upsilon(\mathcal{C})$, the minimum is greater than or equal to b . If the minimum is greater than b , then we can write $a = (a - e_k) + e_k$, where $a - e_k$ is a b -cover and e_k is a 0-cover, a contradiction to the indecomposability of a .

Next we show that $\mathcal{C}^a$ is indecomposable. We proceed by contradiction. Assume that $\mathcal{C}^a$ is decomposable. Then there is a partition X_1, X_2 of X^a such that $\alpha_0(\mathcal{C}^a) = \alpha_0(\mathcal{C}^a[X_1]) + \alpha_0(\mathcal{C}^a[X_2])$. For $1 \leq i \leq n$, we set

$$\ell_i = |X^{a_i} \cap X_1| \quad \text{and} \quad p_i = |X^{a_i} \cap X_2|$$

if $1 \leq i \leq m$ and $\ell_i = p_i = 0$ if $i > m$. Consider the vectors $\ell = (\ell_i)$ and $p = (p_i)$. Notice that a has a decomposition $a = \ell + p$ because one has a partition $X^{a_i} = (X^{a_i} \cap X_1) \cup (X^{a_i} \cap X_2)$ for $1 \leq i \leq m$. To derive a contradiction we now claim that ℓ (resp. p) is an $\alpha_0(\mathcal{C}^a[X_1])$ -cover (resp. $\alpha_0(\mathcal{C}^a[X_2])$ -cover) of $\Upsilon(\mathcal{C})$. Take an arbitrary C in $\Upsilon(\mathcal{C})$. The set

$$C_a = \bigcup_{x_i \in C} \{x_i^1, \dots, x_i^{a_i}\} = \bigcup_{x_i \in C} X^{a_i}$$

is a vertex cover of $\mathcal{C}^a$. Indeed, if f_k is any edge of $\mathcal{C}^a$, then f_k has the form

$$(2.4) \quad f_k = \{x_{k_1}^{j_{k_1}}, x_{k_2}^{j_{k_2}}, \dots, x_{k_r}^{j_{k_r}}\} \quad (1 \leq k_1 < \dots < k_r \leq m; 1 \leq j_{k_i} \leq a_{k_i})$$

for some edge $\{x_{k_1}, x_{k_2}, \dots, x_{k_r}\}$ of $\mathcal{C}$. Since $\{x_{k_1}, x_{k_2}, \dots, x_{k_r}\} \cap C \neq \emptyset$, we get $f_k \cap C_a \neq \emptyset$. Thus C_a is a vertex cover of $\mathcal{C}^a$. Therefore $C_a \cap X_1$ and $C_a \cap X_2$ are vertex covers of $\mathcal{C}^a[X_1]$ and $\mathcal{C}^a[X_2]$ respectively because $E(\mathcal{C}^a[X_i])$ is contained in $E(\mathcal{C}^a)$ for $i = 1, 2$. Hence using the partitions

$$C_a \cap X_1 = \bigcup_{x_i \in C} (X^{a_i} \cap X_1) \quad \text{and} \quad C_a \cap X_2 = \bigcup_{x_i \in C} (X^{a_i} \cap X_2)$$

we obtain

$$\alpha_0(\mathcal{C}^a[X_1]) \leq |C_a \cap X_1| = \sum_{x_i \in C} \ell_i \quad \text{and} \quad \alpha_0(\mathcal{C}^a[X_2]) \leq |C_a \cap X_2| = \sum_{x_i \in C} p_i.$$

This completes the proof of the claim. Consequently a is a decomposable b -cover of $\Upsilon(\mathcal{C})$, where $b = \alpha_0(\mathcal{C}^a)$, a contradiction to the indecomposability of a .

$\Leftarrow$) Assume that $\mathcal{C}^a$ is an indecomposable clutter and $b = \alpha_0(\mathcal{C}^a)$. To show that $x^a t^b$ is a minimal generator of $R_s(I(\mathcal{C}))$ we need only show that a is an indecomposable b -cover of $\Upsilon(\mathcal{C})$. To begin with, notice that a is a b -cover of $\Upsilon(\mathcal{C})$ by Proposition 2.5. We proceed by contradiction assuming that there is a decomposition $a = \ell + p$, where $\ell = (\ell_i)$ is a c -cover of $\Upsilon(\mathcal{C})$, $p = (p_i)$ is a d -cover of $\Upsilon(\mathcal{C})$, and $b = c + d$. Each X^{a_i} can be decomposed as $X^{a_i} = X^{\ell_i} \cup X^{p_i}$, where $X^{\ell_i} \cap X^{p_i} = \emptyset$, $\ell_i = |X^{\ell_i}|$, and $p_i = |X^{p_i}|$. We set

$$X^\ell = X^{\ell_1} \cup \dots \cup X^{\ell_m} \quad \text{and} \quad X^p = X^{p_1} \cup \dots \cup X^{p_m}.$$

Then one has a decomposition $X^a = X^\ell \cup X^p$ of the vertex set of $\mathcal{C}^a$. We now show that $\alpha_0(\mathcal{C}^a[X^\ell]) \geq c$ and $\alpha_0(\mathcal{C}^a[X^p]) \geq d$. By symmetry, it suffices to prove the first inequality. Take an arbitrary minimal vertex cover C_ℓ of $\mathcal{C}^a[X^\ell]$. Then $C_\ell \cup X^p$ is a vertex cover of $\mathcal{C}^a$ because if f is an edge of $\mathcal{C}^a$ contained in X^ℓ , then f is covered by C_ℓ , otherwise f is covered by X^p . Hence there is a minimal vertex cover C_a of $\mathcal{C}^a$ such that $C_a \subset C_\ell \cup X^p$. Since $\mathcal{C}[\{x_1, \dots, x_m\}]$ is a subclutter of $\mathcal{C}^a$, there is a minimal vertex cover C_1 of $\mathcal{C}[\{x_1, \dots, x_m\}]$ contained in C_a . Then

the set $C_1 \cup \{x_i \mid i > m\}$ is a vertex cover of $\mathcal{C}$. Therefore there is a minimal vertex cover C of $\mathcal{C}$ such that $C \cap \{x_1, \dots, x_m\} \subset C_a$. Altogether one has:

$$(2.5) \quad C \cap \{x_1, \dots, x_m\} \subset C_a \subset C_\ell \cup X^p \implies$$

$$(2.6) \quad C \cap \{x_1, \dots, x_m\} \subset C_a \cap \{x_1, \dots, x_m\} \subset (C_\ell \cup X^p) \cap \{x_1, \dots, x_m\}.$$

We may assume that $C_a \cap \{x_1, \dots, x_m\} = \{x_1, \dots, x_s\}$. Next we claim that $X^{a_i} \subset C_a$ for $1 \leq i \leq s$. Take an integer i between 1 and s . Since C_a is a minimal vertex cover of $\mathcal{C}^a$, there exists an edge e of $\mathcal{C}^a$ such that $e \cap C_a = \{x_i^1\}$. Then $(e \setminus \{x_i^1\}) \cup \{x_i^j\}$ is an edge of $\mathcal{C}^a$ for $j = 1, \dots, a_i$, this follows using that the edges of $\mathcal{C}^a$ are of the form described in Eq. (2.4). Consequently $x_i^j \in C_a$ for $j = 1, \dots, a_i$. This completes the proof of the claim. Thus one has $X^{\ell_i} \subset X^{a_i} \subset C_a$ for $1 \leq i \leq s$. Hence, by Eq. (2.5), and noticing that $X^{\ell_i} \cap X^p = \emptyset$, we get $X^{\ell_i} \subset C_\ell$ for $1 \leq i \leq s$. So, using that $\ell_i = 0$ for $i > m$, we get

$$\alpha_0(\mathcal{C}^a[X^\ell]) \geq |C_\ell| \geq \sum_{i=1}^s \ell_i \geq \sum_{x_i \in C \cap \{x_1, \dots, x_m\}} \ell_i = \sum_{x_i \in C} \ell_i \geq c.$$

Therefore $\alpha_0(\mathcal{C}^a[X^\ell]) \geq c$. Similarly $\alpha_0(\mathcal{C}^a[X^p]) \geq d$. Thus

$$\alpha_0(\mathcal{C}^a[X^\ell]) + \alpha_0(\mathcal{C}^a[X^p]) \geq c + d = b,$$

and consequently by Eq. (2.3) we have the equality

$$\alpha_0(\mathcal{C}^a[X^\ell]) + \alpha_0(\mathcal{C}^a[X^p]) = \alpha_0(\mathcal{C}^a).$$

Thus we have shown that $\mathcal{C}^a$ is a decomposable clutter, a contradiction. $\square$

Let $\mathcal{C}$ be a clutter. A set of edges of $\mathcal{C}$ is called *independent* if no two of them have a common vertex. We denote the maximum number of independent edges of $\mathcal{C}$ by $\beta_1(\mathcal{C})$, this number is called the *matching number* of $\mathcal{C}$. In general the vertex covering number and the matching number satisfy $\beta_1(\mathcal{C}) \leq \alpha_0(\mathcal{C})$.

Definition 2.7. If $\beta_1(\mathcal{C}) = \alpha_0(\mathcal{C})$, we say that $\mathcal{C}$ has the *König property*.

Lemma 2.8. *If $\mathcal{C}$ is an indecomposable clutter with the König property, then either $\mathcal{C}$ has no edges and has exactly one isolated vertex or $\mathcal{C}$ has only one edge and no isolated vertices.*

PROOF. Let $f_1, \dots, f_g$ be a set of independent edges and let $X' = \cup_{i=1}^g f_i$, where $g = \alpha_0(\mathcal{C})$. Note that $g = 0$ if $\mathcal{C}$ has no edges. Then $V(\mathcal{C})$ has a partition

$$V(\mathcal{C}) = (\cup_{i=1}^g f_i) \cup (\cup_{x_i \in V(\mathcal{C}) \setminus X'} \{x_i\}).$$

As $\mathcal{C}$ is indecomposable, we get that either $g = 0$ and $V(\mathcal{C}) = \{x_i\}$ for some vertex x_i or $g = 1$ and $V(\mathcal{C}) = f_i$ for some i . Thus in the second case, as $\mathcal{C}$ is a clutter, we get that $\mathcal{C}$ has exactly one edge and no isolated vertices. $\square$

Corollary 2.9. Let $\mathcal{C}$ be a clutter and let $I = I(\mathcal{C})$ be its edge ideal. Then all indecomposable parallelizations of $\mathcal{C}$ satisfy the König property if and only if $I^i = I^{(i)}$ for $i \geq 1$.

PROOF. $\Rightarrow$) It suffices to prove that $R[It] = R_s(I)$. Clearly $R[It] \subset R_s(I)$. To prove the reverse inclusion take a minimal generator $x^a t^b$ of $R_s(I)$. If $b = 0$, then $a = e_i$ for some i and $x^a t^b = x_i$. Thus $x^a t^b \in R[It]$. Assume $b \geq 1$. By Theorem 2.6 we have that $\mathcal{C}^a$ is an indecomposable clutter such that $b = \alpha_0(\mathcal{C}^a)$. As $\mathcal{C}^a$ is indecomposable and satisfies the König property, using Lemma 2.8, it is

not hard to see that $b = 1$ and that $E(\mathcal{C}^a) = \{e\}$ consists of a single edge e of $\mathcal{C}$, i.e., $x^a t^b = x_e t$, where $x_e = \prod_{x_i \in e} x_i$. Thus $x^a t^b \in R[It]$.

$\Leftarrow$) Since $R[It] = R_s(I)$, by Theorem 2.6 we obtain that the only indecomposable parallelizations are either induced subclutters of $\mathcal{C}$ with exactly one edge and no isolated vertices or subclutters consisting of exactly one isolated vertex. Thus in both cases they satisfy the König property. $\square$

A clutter $\mathcal{C}$ is called *Mengerian* if all its parallelizations have the König property. A clutter $\mathcal{C}$ satisfies the *max-flow min-cut property* if the linear program:

$$\max\{\langle \mathbf{1}, y \rangle \mid y \geq 0, Ay \leq a\}$$

has an integral optimal solution for all $a \in \mathbb{N}^n$, where A is the incidence matrix of the clutter $\mathcal{C}$ and $\mathbf{1}$ is the vector of all ones. The columns of A are the characteristic vectors of the edges of $\mathcal{C}$. It is well known that a clutter is Mengerian if and only if it satisfies the max-flow min-cut property [25, Chapter 79].

Thus the last corollary can be restated as:

Corollary 2.10. [17, Corollary 3.14] Let $\mathcal{C}$ be a clutter and let I be its edge ideal. Then $\mathcal{C}$ has the max-flow min-cut property if and only if $I^i = I^{(i)}$ for $i \geq 1$.

The following was the first deep result in the study of symbolic powers of edge ideals from the viewpoint of graph theory.

Corollary 2.11. [28, Theorem 5.9] Let G be a graph and let I be its edge ideal. Then G is bipartite if and only if $I^i = I^{(i)}$ for $i \geq 1$.

PROOF. $\Rightarrow$) If G is a bipartite graph, then any parallelization of G is again a bipartite graph. This means that any parallelization of G satisfies the König property because bipartite graphs satisfy this property [7, Theorem 2.1.1]. Thus $I^i = I^{(i)}$ for all i by Corollary 2.9.

$\Leftarrow$) Assume that $I^i = I^{(i)}$ for $i \geq 1$. By Corollary 2.9 all indecomposable induced subgraphs of G have the König property. If G is not bipartite, then G has an induced odd cycle, a contradiction because induced odd cycles are indecomposable [19] and do not satisfy the König property. $\square$

Corollary 2.12. Let $\mathcal{C}$ be a clutter with vertex set $X = \{x_1, \dots, x_n\}$ and let $S \subset X$. Then the induced clutter $H = \mathcal{C}[S]$ is indecomposable if and only if the monomial $(\prod_{x_i \in S} x_i) t^{\alpha_0(H)}$ is a minimal generator of $R_s(I(\mathcal{C}))$.

PROOF. Let $a = \sum_{x_i \in S} e_i$. Since $\mathcal{C}^a = \mathcal{C}[S]$, the result follows from Theorem 2.6. $\square$

Corollary 2.13. Let $\mathcal{C}$ be a clutter with n vertices and let A be its incidence matrix. If the polyhedron $Q(A) = \{x \mid x \geq 0; xA \geq \mathbf{1}\}$ has only integral vertices, then $\alpha_0(\mathcal{C}^a) \leq n - 1$ for all indecomposable parallelizations $\mathcal{C}^a$ of $\mathcal{C}$.

PROOF. Let $v_1, \dots, v_q$ be the characteristic vectors of the edges of $\mathcal{C}$ and let $\overline{I^i}$ be the integral closure of I^i , where I is the edge ideal of $\mathcal{C}$. As $Q(A)$ is integral, by [17, Corollary 3.13] we have that $\overline{I^i} = I^{(i)}$ for $i \geq 1$, where

$$\overline{I^i} = (\{x^a \mid a \in iB \cap \mathbb{Z}^n\})$$

and $B = \mathbb{Q}_+^n + \text{conv}(v_1, \dots, v_q)$, see [31]. Thus we have the equality $\overline{R[It]} = R_s(I)$, where $\overline{R[It]}$ is the integral closure of $R[It]$ in its field of fractions. Take any

indecomposable parallelization $\mathcal{C}^a$ of $\mathcal{C}$ and consider the monomial $m = x^a t^b$, where $b = \alpha_0(\mathcal{C}^a)$. By Theorem 2.6 m is a minimal generator of $R_s(I)$. Now, according to [14, Corollary 3.11], a minimal generator of $\overline{R[It]}$ has degree in t at most $n - 1$, i.e., $b \leq n - 1$. $\square$

We end this section showing some very basic properties of indecomposable clutters. If e is an edge of a clutter $\mathcal{C}$, we denote by $\mathcal{C} \setminus \{e\}$ the spanning subclutter of $\mathcal{C}$ obtained by deleting e and keeping all the vertices of $\mathcal{C}$.

Definition 2.14. A clutter $\mathcal{C}$ is called *vertex critical* if $\alpha_0(\mathcal{C} \setminus \{x_i\}) < \alpha_0(\mathcal{C})$ for all $x_i \in V(\mathcal{C})$. A clutter $\mathcal{C}$ is called *edge critical* if $\alpha_0(\mathcal{C} \setminus \{e\}) < \alpha_0(\mathcal{C})$ for all $e \in E(\mathcal{C})$.

The next lemma is not hard to prove.

Lemma 2.15. Let x_i be a vertex of a clutter $\mathcal{C}$ and let e be an edge of $\mathcal{C}$.

- (a) If $\alpha_0(\mathcal{C} \setminus \{x_i\}) < \alpha_0(\mathcal{C})$, then $\alpha_0(\mathcal{C} \setminus \{x_i\}) = \alpha_0(\mathcal{C}) - 1$.
- (b) If $\alpha_0(\mathcal{C} \setminus \{e\}) < \alpha_0(\mathcal{C})$, then $\alpha_0(\mathcal{C} \setminus \{e\}) = \alpha_0(\mathcal{C}) - 1$.

Definition 2.16. A clutter $\mathcal{C}$ is called *connected* if there is no $U \subset V(\mathcal{C})$ such that $\emptyset \subsetneq U \subsetneq V(\mathcal{C})$ and such that $e \subset U$ or $e \subset V(\mathcal{C}) \setminus U$ for each edge e of $\mathcal{C}$.

Proposition 2.17. If a clutter $\mathcal{C}$ is indecomposable, then it is connected and vertex critical.

PROOF. Assume that $\mathcal{C}$ is disconnected. Then there is a partition X_1, X_2 of $V(\mathcal{C})$ such that

$$(2.7) \quad E(\mathcal{C}) \subset E(\mathcal{C}[X_1]) \cup E(\mathcal{C}[X_2]).$$

For $i = 1, 2$, let C_i be a minimal vertex cover of $\mathcal{C}[X_i]$ with $\alpha_0(\mathcal{C}[X_i])$ vertices. Then, by Eq. (2.7), $C_1 \cup C_2$ is a minimal vertex cover of $\mathcal{C}$. Hence $\alpha_0(\mathcal{C}[X_1]) + \alpha_0(\mathcal{C}[X_2])$ is greater than or equal to $\alpha_0(\mathcal{C})$. So $\alpha_0(\mathcal{C})$ is equal to $\alpha_0(\mathcal{C}[X_1]) + \alpha_0(\mathcal{C}[X_2])$, a contradiction to the indecomposability of $\mathcal{C}$. Thus $\mathcal{C}$ is connected.

We now show that $\alpha_0(\mathcal{C} \setminus \{x_i\}) < \alpha_0(\mathcal{C})$ for all i . If $\alpha_0(\mathcal{C} \setminus \{x_i\}) = \alpha_0(\mathcal{C})$, then $V(\mathcal{C}) = X_1 \cup X_2$, where $X_1 = V(\mathcal{C}) \setminus \{x_i\}$ and $X_2 = \{x_i\}$. Note that $\mathcal{C}[X_1] = \mathcal{C} \setminus \{x_i\}$. As $\alpha_0(\mathcal{C}[X_1]) = \alpha_0(\mathcal{C})$ and $\alpha_0(\mathcal{C}[X_2]) = 0$, we contradict the indecomposability of $\mathcal{C}$. Thus $\alpha_0(\mathcal{C} \setminus \{x_i\}) < \alpha_0(\mathcal{C})$ and $\mathcal{C}$ is vertex critical. $\square$

Proposition 2.18. If $\mathcal{C}$ is a connected edge critical clutter, then $\mathcal{C}$ is indecomposable.

PROOF. Assume that $\mathcal{C}$ is decomposable. Then there is a partition X_1, X_2 of $V(\mathcal{C})$ into nonempty vertex sets such that $\alpha_0(\mathcal{C}) = \alpha_0(\mathcal{C}[X_1]) + \alpha_0(\mathcal{C}[X_2])$. Since $\mathcal{C}$ is connected, there is an edge $e \in E(\mathcal{C})$ intersecting both X_1 and X_2 . Pick a minimal vertex cover C of $\mathcal{C} \setminus \{e\}$ with less than $\alpha_0(\mathcal{C})$ vertices. As $E(\mathcal{C}[X_i])$ is a subset of $E(\mathcal{C} \setminus \{e\}) = E(\mathcal{C}) \setminus \{e\}$ for $i = 1, 2$, we get that C covers all edges of $\mathcal{C}[X_i]$ for $i = 1, 2$. Hence C must have at least $\alpha_0(\mathcal{C})$ vertices, a contradiction. $\square$

From Propositions 2.17 and 2.18 we obtain:

Corollary 2.19. The following hold for any connected clutter :

$$\text{edge critical} \implies \text{indecomposable} \implies \text{vertex critical}.$$

The next result can be used to build indecomposable clutters.

Proposition 2.20. Let $\mathcal{D}$ be a clutter obtained from a clutter $\mathcal{C}$ by adding a new vertex v and some new edges containing v and some vertices of $V(\mathcal{C})$. If $a = (1, \dots, 1) \in \mathbb{N}^n$ is an indecomposable $\alpha_0(\mathcal{C})$ -cover of $\Upsilon(\mathcal{C})$ such that $\alpha_0(\mathcal{D}) = \alpha_0(\mathcal{C}) + 1$, then $a' = (a, 1)$ is an indecomposable $\alpha_0(\mathcal{D})$ -cover of $\Upsilon(\mathcal{D})$.

PROOF. Clearly a' is an $\alpha_0(\mathcal{D})$ -cover of $\Upsilon(\mathcal{D})$. Assume that $a' = a'_1 + a'_2$, where $a'_i \neq 0$ is a b'_i -cover of $\Upsilon(\mathcal{D})$ and $b'_1 + b'_2 = \alpha_0(\mathcal{D})$. We may assume that $a'_1 = (1, \dots, 1, 0, \dots, 0)$ and $a'_2 = (0, \dots, 0, 1, \dots, 1)$. Let a_i be the vector in $\mathbb{N}^n$ obtained from a'_i by removing its last entry. Set $v = x_{n+1}$. Take a minimal vertex cover C_k of $\mathcal{C}$ and consider $C'_k = C_k \cup \{x_{n+1}\}$. Let u'_k (resp. u_k) be the characteristic vector of C'_k (resp. C_k). Then

$$\langle a_1, u_k \rangle = \langle a'_1, u'_k \rangle \geq b'_1 \text{ and } \langle a_2, u_k \rangle + 1 = \langle a'_2, u'_k \rangle \geq b'_2,$$

and consequently a_1 is a b'_1 -cover of $\Upsilon(\mathcal{C})$. If $b'_2 = 0$, then a_1 is an $\alpha_0(\mathcal{D})$ -cover of $\Upsilon(\mathcal{C})$, a contradiction; because if u is the characteristic vector of a minimal vertex cover of $\mathcal{C}$ with $\alpha_0(\mathcal{C})$ elements, then we would obtain $\alpha_0(\mathcal{C}) \geq \langle u, a_1 \rangle \geq \alpha_0(\mathcal{D})$, which is impossible. Thus $b'_2 \geq 1$, and a_2 is a $(b'_2 - 1)$ -cover of $\Upsilon(\mathcal{C})$ if $a_2 \neq 0$. Hence $a_2 = 0$, because $a = a_1 + a_2$ and a is indecomposable. This means that $a'_2 = e_{n+1}$ is a b'_2 -cover of $\Upsilon(\mathcal{D})$, a contradiction. Therefore a' is an indecomposable $\alpha_0(\mathcal{D})$ -cover of $\Upsilon(\mathcal{D})$, as required. $\square$

3. Indecomposable parallelizations and Hilbert bases

Let $\mathcal{C}$ be a clutter with vertex set $X = \{x_1, \dots, x_n\}$ and let $C_1, \dots, C_s$ be the minimal vertex covers of $\mathcal{C}$. For $1 \leq k \leq n$, we denote the characteristic vector of C_k by u_k .

The *Simis cone* of $I = I(\mathcal{C})$ is the rational polyhedral cone:

$$\text{Cn}(I) = H_{e_1}^+ \cap \dots \cap H_{e_{n+1}}^+ \cap H_{(u_1, -1)}^+ \cap \dots \cap H_{(u_s, -1)}^+.$$

Here H_a^+ denotes the closed halfspace $H_a^+ = \{x \mid \langle x, a \rangle \geq 0\}$ and H_a stands for the hyperplane through the origin with normal vector a . Simis cones were introduced in [14] to study symbolic Rees algebras of square-free monomial ideals. The term *Simis cone* is intended to do homage to Aron Simis [26, 27, 28]. The Simis cone is a pointed rational polyhedral cone. By [24, Theorem 16.4] there is a unique minimal finite set of integral vectors

$$\mathcal{H} = \{h_1, \dots, h_r\} \subset \mathbb{Z}^{n+1}$$

such that $\mathbb{Z}^{n+1} \cap \mathbb{R}_+ \mathcal{H} = \mathbb{N} \mathcal{H}$ and $\text{Cn}(I) = \mathbb{R}_+ \mathcal{H}$ (minimal relative to taking subsets), where $\mathbb{R}_+ \mathcal{H}$ denotes the cone generated by $\mathcal{H}$ consisting of all linear combinations of $\mathcal{H}$ with non-negative real coefficients and $\mathbb{N} \mathcal{H}$ denotes the semigroup generated by $\mathcal{H}$ consisting of all linear combinations of $\mathcal{H}$ with coefficients in $\mathbb{N}$. The set $\mathcal{H}$ is called the *Hilbert basis* of $\text{Cn}(I)$. The Hilbert basis of $\text{Cn}(I)$ has the following useful description.

Theorem 3.1. [24, p. 233] $\mathcal{H}$ is the set of all integral vectors $0 \neq h \in \text{Cn}(I)$ such that h is not the sum of two other non-zero integral vectors in $\text{Cn}(I)$.

Corollary 3.2. Let $\mathcal{H}$ be the Hilbert basis of $\text{Cn}(I)$. Then

$$(3.1) \quad \mathcal{H} = \{(a, b) \mid x^a t^b \text{ is a minimal generator of } R_s(I)\}$$

$$(3.2) \quad = \{(a, \alpha_0(\mathcal{C})) \mid \mathcal{C}^a \text{ is an indecomposable parallelization of } \mathcal{C}\}$$

and $R_s(I)$ is equal to the semigroup ring $K[\mathbb{N} \mathcal{H}]$ of $\mathbb{N} \mathcal{H}$.

PROOF. The first equality follows from Lemma 2.1 and Theorem 3.1. The second equality follows from Theorem 2.6. The equality $K[\mathbb{N}\mathcal{H}] = \mathbb{N}\mathcal{H}$ was first observed in [14, Theorem 3.5]. $\square$

This result is interesting because it allows to compute all indecomposable parallelizations of $\mathcal{C}$ and all indecomposable induced subclutters of $\mathcal{C}$ using Hilbert bases. In particular, as is seen in Corollary 3.3, we can use this result to decide whether any given graph or clutter is indecomposable (see Example 3.4).

The indecomposable subclutters can be computed using the next consequence of Corollary 3.2.

Corollary 3.3. Let $\mathcal{C}$ be a clutter and let $\alpha = (a_1, \dots, a_n, b)$ be a vector in $\{0, 1\}^n \times \mathbb{N}$. Then α is in the Hilbert basis of $\text{Cn}(I(\mathcal{C}))$ if and only if the induced subclutter $H = \mathcal{C}[\{x_i \mid a_i = 1\}]$ is indecomposable with $b = \alpha_0(H)$.

Example 3.4. Consider the graph G shown below. Let I be the edge ideal of G and let $\mathcal{H}$ be the Hilbert basis of $\text{Cn}(I)$. Using Corollary 3.2, together with NORMALIZ [4], it is seen that G has exactly 61 indecomposable parallelizations and 49 indecomposable subgraphs. Since $\alpha_0(G) = 6$ and the vector $(1, \dots, 1, 6)$ is not in $\mathcal{H}$ we obtain that G is a decomposable graph.




Fig. 4. Decomposable graph G

The vector $a = (1, \dots, 1, 2, 7)$ is in $\mathcal{H}$, i.e., $G^{(1, \dots, 1, 2)}$ is indecomposable and has covering number 7.




Fig. 5. Indecomposable graph $G^{(1, \dots, 1, 2)}$

The next result, together with Corollary 3.3, allows to locate all *induced odd cycles* (odd holes) and all *induced complements of odd cycles* (odd antiholes).

Lemma 3.5. Let $C_n = \{x_1, \dots, x_n\}$ be a cycle. (a) If $n \geq 5$ is odd, then the complement C'_n of C_n is an indecomposable graph, (b) if n is odd, then C_n is an indecomposable cycle, and (c) any complete graph is indecomposable.

PROOF. (a) Assume that $G = C'_n$ is decomposable. Then there are disjoint sets X_1, X_2 such that $V(G) = X_1 \cup X_2$ and $\alpha_0(G) = \alpha_0(G[X_1]) + \alpha_0(G[X_2])$. Since $\beta_0(G) = 2$, it is seen that $G[X_i]$ is a complete graph for $i = 1, 2$. We may assume that $x_1 \in X_1$. Then x_2 must be in X_2 , otherwise $\{x_1, x_2\}$ is an

edge of $G[X_1]$, a contradiction. By induction it follows that $x_1, x_3, x_5, \dots, x_n$ are in X_1 . Consequently $\{x_1, x_n\}$ is an edge of $G[X_1]$, a contradiction. Thus G is indecomposable. (b) This was observed in [19]. (c) Follows readily from the fact that the covering number of a complete graph in r vertices is $r - 1$. $\square$

Example 3.6. Consider the graph G of Fig. 6, where vertices are labeled with i instead of x_i . Using Corollary 3.2, together with NORMALIZ [4], it is seen that G has exactly 21 indecomposable parallelizations, 20 of which correspond to indecomposable subgraphs. Apart from the seven vertices, the nine edges, one triangle and three pentagons, the only indecomposable parallelization of G which is not a subgraph is the duplication shown in Fig 7.




Fig. 6. Decomposable graph G




Fig. 7. Indecomposable graph $G^{(2,1,1,1,1,1,1)}$

Example 3.7. Consider the graph G of Fig. 8. Using Corollary 3.2 and NORMALIZ [4], it is seen that G has exactly 103 indecomposable parallelizations, 92 of which correspond to indecomposable subgraphs. The only indecomposable parallelization G^a which do not delete vertices is that obtained by duplication of the five outer vertices, i.e., $a = (2, 2, 2, 2, 2, 1, 1, 1, 1, 1)$ and $\alpha_0(G^a) = 11$.




Fig. 8. Decomposable graph G

4. Symbolic Rees algebras and perfect graphs

We now turn our attention to the indecomposability of graphs and its connection with the theory of perfect graphs. Examples of indecomposable graphs include complete graphs, odd cycles, and complements of odd cycles of length at least 5 (see Lemma 3.5).

Let us recall the notion of a perfect graph that was introduced by Berge [2, Chapter 16]. A *colouring* of the vertices of a graph G is an assignment of colours to the vertices of G in such a way that adjacent vertices have distinct colours. The *chromatic number* of G is the minimal number of colours in a colouring of G . A graph is *perfect* if for every induced subgraph H , the chromatic number of H

equals the size of the largest complete subgraph of H . We refer to [6, 18, 25] for the theory of perfect graphs.

The next result shows that indecomposable graphs occur naturally in the theory of perfect graphs.

Proposition 4.1. [12, Proposition 2.13] A graph G is perfect if and only if the indecomposable parallelizations of G are exactly the complete subgraphs of G

Let G be a graph. We denote a complete subgraph of G with r vertices by $\mathcal{K}_r$. The empty set is regarded as an independent set of vertices whose characteristic vector is the zero vector. A *clique* of G is a subset of the set of vertices that induces a complete subgraph. The *support* of a monomial $x^a = x_1^{a_1} \cdots x_n^{a_n}$, denoted by $\text{supp}(x^a)$, is the set $\text{supp}(x^a) = \{x_i \mid a_i > 0\}$. If $a_i \in \{0, 1\}$ for all i , x^a is called a *square-free* monomial.

The next major result shows that the symbolic Rees algebra of the edge ideal of a perfect graph G is completely determined by the cliques of G . This was first shown in [32] using polyhedral geometry.

Corollary 4.2. [32, Corollary 3.3] If G is a perfect graph, then

$$R_s(I(G)) = K[\{x^a t^b \mid x^a \text{ is square-free ; } G[\text{supp}(x^a)] = \mathcal{K}_{b+1}\}].$$

PROOF. Let $x^a t^b$ be a minimal of $R_s(I(G))$. By Theorem 2.6 G^a is an indecomposable graph and $b = \alpha_0(G^a)$. As G is perfect, by Proposition 4.1, we obtain that G^a is a complete subgraph of G with $b + 1$ vertices. $\square$

Since complete graphs are perfect, an immediate consequence is:

Corollary 4.3. [1] If G is a complete graph, then

$$R_s(I(G)) = K[\{x^a t^b \mid x^a \text{ is square-free ; } \deg(x^a) = b + 1\}].$$

References

1. C. Bahiano, Symbolic powers of edge ideals, J. Algebra **273** (2004), no. 2, 517–537.
2. C. Berge, *Graphs and hypergraphs*, North-Holland Mathematical Library, Vol. 6, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1976.
3. J. P. Brennan, L. A. Dupont and R. H. Villarreal, Duality, a-invariants and canonical modules of rings arising from linear optimization problems, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **51** (2008), no. 4, 279–305.
4. W. Bruns and B. Ichim, NORMALIZ 2.0, Computing normalizations of affine semigroups 2008. Available from <http://www.math.uos.de/normaliz>.
5. M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem, Ann. of Math. (2) **164** (2006), no. 1, 51–229.
6. G. Cornuéjols, *Combinatorial optimization: Packing and covering*, CBMS-NSF Regional Conference Series in Applied Mathematics **74**, SIAM (2001).
7. R. Diestel, *Graph Theory*, Graduate Texts in Mathematics **173**, Springer-Verlag, New York, 2nd ed., 2000.
8. L. A. Dupont, C. Rentería and R. H. Villarreal, Systems with the integer rounding property in normal monomial subrings, An. Acad. Brasil. Ciênc, to appear.
9. L. A. Dupont, E. Reyes and R. H. Villarreal, Cohen-Macaulay clutters with combinatorial optimization properties and parallelizations of normal edge ideals, São Paulo J. Math. Sci. **3** (2009), no. 1, 61–75.
10. L. A. Dupont and R. H. Villarreal, Edge ideals of clique clutters of comparability graphs and the normality of monomial ideals, Math. Scand. **106** (2010), no. 1, 88–98.
11. L. A. Dupont and R. H. Villarreal, Algebraic and combinatorial properties of ideals and algebras of uniform clutters of TDI systems, J. Comb. Optim., to appear.

12. L. A. Dupont and R. H. Villarreal, Symbolic Rees algebras, vertex covers and irreducible representations of Rees cones, *Algebra Discrete Math.*, to appear.
13. P. Erdős and T. Gallai, On the minimal number of vertices representing the edges of a graph, *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **6** (1961), 181–203.
14. C. Escobar, R. H. Villarreal and Y. Yoshino, Torsion freeness and normality of blowup rings of monomial ideals, *Commutative Algebra*, Lect. Notes Pure Appl. Math. **244**, Chapman & Hall/CRC, Boca Raton, FL, 2006, pp. 69–84.
15. C. A. Francisco, H. T. Hà and A. Van Tuyl, Associated primes of monomial ideals and odd holes in graphs, *J. Algebraic Combin.* **32** (2010), no. 2, 287–301.
16. I. Gitler, E. Reyes and R. H. Villarreal, Blowup algebras of square-free monomial ideals and some links to combinatorial optimization problems, *Rocky Mountain J. Math.* **39** (2009), no. 1, 71–102.
17. I. Gitler, C. Valencia and R. H. Villarreal, A note on Rees algebras and the MFMC property, *Beiträge Algebra Geom.* **48** (2007), No. 1, 141–150.
18. M. C. Golumbic, Algorithmic graph theory and perfect graphs, second edition, *Annals of Discrete Mathematics* **57**, Elsevier Science B.V., Amsterdam, 2004.
19. F. Harary and M. D. Plummer, On indecomposable graphs, *Canad. J. Math.* **19** (1967), 800–809.
20. J. Herzog, T. Hibi and N. V. Trung, Symbolic powers of monomial ideals and vertex cover algebras, *Adv. Math.* **210** (2007), 304–322.
21. G. Lyubeznik, On the arithmetical rank of monomial ideals, *J. Algebra* **112** (1988), 86–89.
22. J. Martínez-Bernal, E. O’Shea and R. H. Villarreal, Ehrhart clutters: regularity and max-flow min-cut, *Electron. J. Combin.* **17** (2010), no. 1, R52.
23. A. Schrijver, On total dual integrality, *Linear Algebra Appl.* **38** (1981), 27–32.
24. A. Schrijver, *Theory of Linear and Integer Programming*, John Wiley & Sons, New York, 1986.
25. A. Schrijver, *Combinatorial Optimization*, Algorithms and Combinatorics **24**, Springer-Verlag, Berlin, 2003.
26. A. Simis, *Combinatoria Algebraica*, XVIII Coloquio Brasileiro de Matematica, IMPA, 1991 (Apendice. Palimpsesto 2: Potencias simbolicas, 2.1).
27. A. Simis and B. Ulrich, On the ideal of an embedded join, *J. Algebra* **226** (2000), no. 1, 1–14.
28. A. Simis, W. V. Vasconcelos and R. H. Villarreal, On the ideal theory of graphs, *J. Algebra*, **167** (1994), 389–416.
29. S. Sullivant, Combinatorial symbolic powers, *J. Algebra* **319** (2008), no. 1, 115–142.
30. W. V. Vasconcelos, *Arithmetic of Blowup Algebras*, London Math. Soc., Lecture Note Series **195**, Cambridge University Press, Cambridge, 1994.
31. R. H. Villarreal, *Monomial Algebras*, Dekker, New York, N.Y., 2001.
32. R. H. Villarreal, Rees algebras and polyhedral cones of ideals of vertex covers of perfect graphs, *J. Algebraic Combin.* **27** (2008), no. 3, 293–305.

DEPARTAMENTO DE MATEMÁTICAS, CENTRO DE INVESTIGACIÓN Y DE ESTUDIOS AVANZADOS DEL IPN, APARTADO POSTAL 14-740, 07000 MEXICO CITY, D.F.

E-mail address: jmb@math.cinvestav.mx

DEPARTAMENTO DE MATEMÁTICAS, ESCUELA SUPERIOR DE FÍSICA Y MATEMÁTICAS, INSTITUTO POLITÉCNICO NACIONAL, 07300 MEXICO CITY, D.F.

E-mail address: renteri@esfm.ipn.mx

DEPARTAMENTO DE MATEMÁTICAS, CENTRO DE INVESTIGACIÓN Y DE ESTUDIOS AVANZADOS DEL IPN, APARTADO POSTAL 14-740, 07000 MEXICO CITY, D.F.

E-mail address: vila@math.cinvestav.mx